

QMW-93-25
October 1993

GAUGED HETEROTIC SIGMA-MODELS

C. M. HULL

*Physics Department, Queen Mary and Westfield College,
Mile End Road, London E1 4NS, United Kingdom.*

ABSTRACT

The gauging of isometries in general sigma-models which include fermionic terms which represent the interaction of strings with background Yang-Mills fields is considered. Gauging is possible only if certain obstructions are absent. The quantum gauge anomaly is discussed, and the (1,0) supersymmetric generalisation of the gauged action given.

Non-linear sigma-models are important two-dimensional field theories and those that are conformally invariant describe the propagation of a string in a curved space-time [1]. Gauging such sigma-models can give a construction of new conformal field theories, and gauged Wess-Zumino-Witten models provide a lagrangian formulation of the coset construction [2]. More recently, duality symmetry in string theory has been formulated in terms of gauged sigma-models [3], and this has led to the proposal of non-abelian generalisations of duality symmetry [4]. Supersymmetric gauged sigma-models have also been used to construct (p,q) supersymmetric integrable models [5]. In [6,7,8], the gauging of general non-linear sigma-models with Wess-Zumino terms was shown to be possible only if certain obstructions were absent, and the gauged action was given. The (p,q) supersymmetric generalisations of these sigma-models were constructed in [9]. The purpose of this paper is to extend these results to the case of sigma-models with fermionic terms, representing the interaction of strings with background Yang-Mills fields C_i^{AB} , in addition to the metric g_{ij} , anti-symmetric tensor gauge field b_{ij} and dilaton Φ . In particular, the (1,0) supersymmetric version of such terms describes the propagation of heterotic strings in backgrounds with non-trivial gauge fields [10] and the gauged version can be used to formulate the effect of duality transformations on Yang-Mills fields [12].

The action for a bosonic two dimensional sigma-model with Wess-Zumino term and Fradkin-Tseytlin term is

$$S_0 = \frac{1}{2} \int d^2x \sqrt{h} (g_{ij} \partial^\mu \phi^i \partial_\mu \phi^j + \epsilon^{\mu\nu} b_{ij} \partial_\mu \phi^i \partial_\nu \phi^j + \Phi R) \quad (1)$$

where the D -dimensional target space M has metric $g_{ij}(\phi)$, coordinates ϕ^i ($i = 1, \dots, D$) and torsion three-form H given in terms of a potential b_{ij} by $H_{ijk} = \frac{3}{2} \partial_{[i} b_{jk]}$. It is invariant under the transformation

$$\delta \phi^i = e \lambda^a \xi_a^i \quad (2)$$

where λ^a ($a = 1, \dots, n$) are infinitesimal constant parameters, e is a constant and ξ_a

are a set of vector fields on M provided the Lie derivative with respect to ξ_a of g_{ij} , Φ and H_{ijk} vanish, which will be the case if

$$\nabla_{(i}\xi_{j)a} = 0 \quad (3)$$

(where $\xi_{ia} = g_{ij}\xi_a^j$)

$$\xi_a^i \partial_i \Phi = 0 \quad (4)$$

and $\xi_a^i H_{ijk}$ is an exact two-form, *i.e.* there is some (globally defined) set of Lie-algebra-valued one-forms v_a , defined up to the addition of a closed form, with components v_{ia} such that

$$\xi_a^i H_{ijk} = \partial_{[j} v_{k]a} \quad (5)$$

Then ξ_a^i are Killing vectors which can be taken to generate some n -dimensional isometry group G satisfying

$$[\xi_a, \xi_b] \equiv \mathcal{L}_a \xi_b = f_{ab}^c \xi_c \quad (6)$$

where f_{ab}^c are the structure constants of G and $\mathcal{L}_a$ denotes the Lie derivative with respect to ξ_a .

The vanishing of the Lie derivative of H_{ijk} implies that

$$\mathcal{L}_a b_{ij} = \partial_{[i} \Lambda_{j]a} \quad (7)$$

for some Λ_{ia} so that the variation of b_{ij} can be cancelled by an anti-symmetric tensor gauge transformation, or equivalently, the variation of the b -term in (1) is a total derivative. If $\Lambda_{ia} = 0$, then the symmetry can be gauged by minimal coupling, *i.e.* by replacing the derivatives ∂_μ in (1) by gauge-covariant derivatives

D_μ , where

$$D_\mu \phi^i = \partial_\mu \phi^i - e A_\mu^a \xi_a^i \quad (8)$$

and the gauge field A_μ^a transforms as

$$\delta A_\mu^a = \partial_\mu \lambda^a + e f_{bc}^a A_\mu^b \lambda^c \quad (9)$$

In the case in which $\Lambda_{ia} \neq 0$, minimal coupling is not sufficient and the gauging is as given in [6,7,8]. The gauging can in principle be given in terms of the Λ_{ia} given in (7), but these are not vector fields in general since the b_{ij} are not tensors but are connections, and it is more convenient to work in terms of the covariant v_{ia} . Gauging of the isometry symmetry (2) is possible only if [6,7,8] (i) the v_{ia} can be chosen to be equivariant, *i.e.* chosen so that

$$\mathcal{L}_a v_{ib} = f_{ab}^c v_{ic} \quad (10)$$

and (ii) if

$$c_{(ab)} = 0 \quad (11)$$

where

$$c_{ab} = v_{ia} \xi_b^i, \quad (12)$$

If these two conditions are satisfied, then the gauged action is

$$\begin{aligned} S_G = & \frac{1}{2} \int d^2 x \sqrt{h} \{ g_{ij} D_\mu \phi^i D^\mu \phi^j + \Phi R \} \\ & + \frac{1}{2} \int d^2 x \sqrt{h} \epsilon^{\mu\nu} \left(b_{ij} \partial_\mu \phi^i \partial_\nu \phi^j + 2e A_\mu^a v_{ia} \partial_\nu \phi^i - e^2 c_{[ab]}(\phi) A_\mu^a A_\nu^b \right) \end{aligned} \quad (13)$$

This can be rewritten as [6,8]

$$\begin{aligned} S_G = & \frac{1}{2} \int d^2 x \sqrt{h} \{ g_{ij} D_\mu \phi^i D^\mu \phi^j + \Phi R \} \\ & + \int_Y \left(\frac{1}{3} H_{ijk} D\phi^i D\phi^j D\phi^k + \frac{e}{2} v_{ia} D\phi^i F^a \right) \end{aligned} \quad (14)$$

where Y is a three-manifold whose boundary is the world-sheet X and the field

strength two-form is $F^a = dA^a - \frac{1}{2}ef_{bc}^a A^b A^c$. As usual, the fields ϕ^i, A_μ^a on X are extended to fields on Y in the second term in (14). If (10) is satisfied but (11) is not, then the action (13) is not gauge-invariant, but satisfies

$$\delta S = e^2 \int d^2x \sqrt{h} \epsilon^{\mu\nu} c_{(ab)} A_\mu^a (\partial_\nu \lambda^b) \quad (15)$$

which is proportional to the consistent chiral anomaly in two dimensions.

Suppose now that one adds a fermionic term of the form [10]

$$S_f = \frac{i}{2} \int d^2x \sqrt{h} \psi_A (\nabla_+ \psi)_A \quad (16)$$

where ψ_A are chiral Majorana world-sheet spinor fields that are also sections of an $O(N)$ vector bundle over M with connection $C_i^{AB}(\phi)$, $C_i^{AB} = -C_i^{BA}$, and fibre metric δ_{AB} , which is used to raise and lower the $O(N)$ vector indices $A, B, \dots = 1, \dots, N$. The covariant derivative is

$$(\nabla_\mu \psi)_A = \partial_\mu \psi_A - \frac{1}{2} \omega_\mu \psi_A - \partial_\mu \phi^i C_i^{AB} \psi_B \quad (17)$$

where $\omega_\mu = \frac{1}{2} \epsilon_{ab} \omega_\mu^{ab}$, ω_μ^{ab} is the world-sheet spin-connection and $\nabla_\pm = e_\pm^\mu \nabla_\mu$ where e_a^μ ($a = \pm$) are zweibeins, with $e_\pm^\mu = \pm \epsilon^{\mu\nu} e_{\nu\pm}$. This is formally invariant under the $O(N)$ gauge transformations

$$\delta \psi_A = M_A^B \psi_B, \quad \delta C_i = \partial_i M - [C_i, M] \quad (18)$$

with parameter $M_A^B(\phi)$. Under an arbitrary variation of the fields, the action (16) changes by

$$\delta S_f = i \int d^2x \sqrt{h} \left\{ (\Delta \psi_A) (\nabla_+ \psi)_A - \frac{1}{2} \delta \phi^i \partial_+ \phi^j G_{ij}^{AB} \psi_A \psi_B \right\} \quad (19)$$

where the field strength is

$$G_{ij} = \partial_i C_j - \partial_j C_i - [C_i, C_j] \quad (20)$$

and the covariant variation is defined by

$$\Delta\psi_A = \delta\psi_A - \delta\phi^i C_i^{AB} \psi_B \quad (21)$$

The transformation (2) for constant λ will lead to a symmetry of the action S_f if the connection C_i is invariant up to a gauge transformation:

$$\mathcal{L}_a C_i = \nabla_i \kappa_a \quad (22)$$

for some $\kappa_a^{AB}(\phi)$, as the variation of the action can then be cancelled by an $O(N)$ transformation of ψ_A , $\delta\psi_A = \lambda^a \kappa_a^{AB} \psi_B$. This condition has been discussed in [11], where particular attention is paid to global aspects. It can be reformulated covariantly as follows. The condition for there to be an isometry symmetry is that the field strength satisfy

$$\xi_a^i G_{ij}^{AB} = \nabla_i \mu_a^{AB} \quad (23)$$

for some $\mu_a^{AB}(\phi)$. This is equivalent to (22) with

$$\mu_a = \kappa_a - \xi_a^i C_i \quad (24)$$

The κ_a are not $O(N)$ -covariant (they transform as a connection), and it is more convenient to work with the μ_a^{AB} defined by (23), which transform covariantly under $O(N)$ transformations

$$\delta\mu_a = [M, \mu_a] \quad (25)$$

just as for the Wess-Zumino term it was better to work with the v_{ia} rather than the Λ_{ia} . If (23) is satisfied, then the action (16) is invariant under the rigid transformations given by (2) and

$$\Delta\psi_A = e\lambda^a \mu_a^{AB} \psi_B \quad (26)$$

Under the transformations given by (2),(26), with local $\lambda(x)$, the action S_f

varies by

$$\delta S_f = e \int d^2x \sqrt{h} \partial_+ \lambda^a J_{a-} \quad (27)$$

where J_{a-} is the Noether current

$$J_{a-} = \frac{i}{2} (\psi_A \mu_a^{AB} \psi_B) \quad (28)$$

This variation can be cancelled by adding the Noether coupling $-e A_+ J_-$ to obtain

$$\begin{aligned} S_g &= \frac{i}{2} \int d^2x \sqrt{h} (\psi_A (\nabla_+ \psi)_A - e A_+^a (\psi \mu_a \psi)) \\ &= \frac{i}{2} \int d^2x \sqrt{h} \psi_A (\mathcal{D}_+ \psi)_A \end{aligned} \quad (29)$$

where

$$\begin{aligned} (\mathcal{D}_+ \psi)_A &= (\nabla_+ \psi)_A - e A_+^a \mu_a^{AB} \psi_B = \partial_+ \psi_A - \mathcal{A}_+^{AB} \psi_B, \\ \mathcal{A}_\mu^{AB} &= e A_\mu^a \mu_a^{AB} + C_i^{AB} \partial_\mu \phi^i \end{aligned} \quad (30)$$

This action is then fully gauge-invariant provided the μ_a are equivariant, *i.e.* they satisfy

$$\hat{\mathcal{L}}_a \mu_b - [\mu_a, \mu_b] = f_{ab}^c \mu_c \quad (31)$$

where $\hat{\mathcal{L}}_a$ is the gauge-covariant Lie derivative, which for tensors $T_{ij\dots}^{AB}$ transforming according to the adjoint of $O(N)$ is given by

$$\hat{\mathcal{L}}_a T_{ij\dots}^{AB} = \mathcal{L}_a T_{ij\dots}^{AB} - \xi_a^i [C_i, T_{ij\dots}]^{AB} \quad (32)$$

This is the analogue of the equivariance condition (10).

To summarise, the action (16) is invariant under rigid isometries provided that the field-strength satisfies (23) for some μ_a , and this can be promoted to a local

symmetry provided the μ_a satisfy the equivariance condition (31), in which case the gauged action is (29). Note that (23) is equivalent to the condition that

$$\hat{\mathcal{L}}_a G_{ij} = [\mu_a, G_{ij}] \quad (33)$$

for some μ_a , so that the gauge-covariant Lie derivative of the field-strength vanishes up to a gauge transformation. The action (29) is also invariant under the $O(N)$ transformations (18),(25).

As an example, consider the case in which the vector bundle is the tangent bundle and C_i^{AB} is the torsion-free spin-connection, where A, B are now tangent space indices with respect to a target space vielbein E_i^A . Then $G_{ij}^{AB} = R_{ijAB}$ where R_{ijAB} is the curvature tensor and (23) is automatically satisfied with μ_a^{AB} proportional to $E_i^A E_j^B \nabla^i \xi_a^j$.

The gauge variation of the connection $\mathcal{A}_+$ defined by (30) is

$$\delta \mathcal{A}_+^{AB} = \partial_+ \lambda^{AB} - [\mathcal{A}, \lambda]^{AB}, \quad \lambda^{AB} \equiv \lambda^a \kappa_a^{AB} \quad (34)$$

so that the action (29) is manifestly invariant under the transformations (34) and $\delta \psi_A = -e \lambda^{AB} \psi_B$, which is equivalent to (26). Quantum mechanically, the gauge symmetry of the chiral fermion action (29) is anomalous, with the variation of the effective action proportional to

$$\Delta = \int d^2 x \sqrt{h} \mathcal{A}_+^{AB} \partial_- \lambda^{AB} \quad (35)$$

Adding a counterterm proportional to $\text{Tr}(\mathcal{A}_+ \mathcal{A}_-)$, the anomaly becomes proportional to

$$\Delta = \int \mathcal{A}^{AB} d\lambda^{AB} \quad (36)$$

where $\mathcal{A}^{AB} = \mathcal{A}_\mu^{AB} dx^\mu$. When rewritten in terms of A^a and λ^a , this contains $d\phi$ and $d\phi d\phi$ terms.

It is straightforward to extend these results to the (1,0) supersymmetric sigma-model. For a flat world-sheet with $h_{\mu\nu} = \eta_{\mu\nu}$, flat (1,0) superspace has coordinates x^+, x^- and θ and flat superspace derivative D with $D^2 = i\partial_+$. The (1,0) supersymmetric generalisation of (1) is given by [10]

$$S = -i \left[\int d^2x d\theta g_{ij} D\phi^i \partial_- \phi^j + \int d^2x dt d\theta H_{ijk} \partial_t \phi^i D\phi^j \partial_- \phi^k \right], \quad (37)$$

where ϕ is now a superfield, *i.e.* a map from the (1,0) superspace into M . The transformations (2), now involving superfields, are rigid symmetries of the action (37) provided that the vector fields ξ_a^i satisfy the same conditions as in the bosonic case. To promote these rigid symmetries to local ones, it is necessary to introduce a (1,0) super Yang-Mills multiplet. This is described by gauge superfields $A(x, \theta)$, $A_+(x, \theta)$, $A_-(x, \theta)$ which can be used to define gauge-covariant derivatives $\nabla, \nabla_-, \nabla_+$. These are taken to satisfy the constraints [13]

$$[\nabla, \nabla] = 2i\nabla_+, \quad [\nabla, \nabla_-] = W, \quad [\nabla_+, \nabla_-] = F, \quad (38)$$

with all other super-commutators equal to zero. In equation (38), the field strength W^a is an unconstrained superfield while the Bianchi identities imply that F^a is proportional to DW . Acting on sigma-model fields, these become (setting $e = -1$)

$$\nabla \phi^i = D\phi^i + A^a \xi_a^i(\phi), \quad [\nabla, \nabla_-] \phi^i = W^a \xi_a^i(\phi) \quad (39)$$

etc. Gauging is possible if and only if it is possible for the corresponding bosonic model, in which case the action for the gauged (1,0) sigma model is

$$S = -i \int d^2x d\theta \left[g_{ij} \nabla \phi^i \nabla_- \phi^j + b_{ij} D\phi^i \partial_- \phi^j - A^a u_{ia} \partial_- \phi^i + A_-^a u_{ia} D\phi^i + A^a A_-^b c_{[ab]} \right], \quad (40)$$

The (1,0) supersymmetric generalisation of (16) is [10]

$$S = \frac{1}{2} \int d^2x d\theta \psi_A \tilde{D} \psi_A \quad (41)$$

where

$$\tilde{D}\psi_A = D\psi_A - D\phi^i C_i^{AB}\psi_B \quad (42)$$

and ψ_A are now fermionic superfields. Again, this is invariant under rigid isometry symmetries if and only if the connection $C_i(\phi)$ satisfies (23) for some μ_a , and the rigid symmetry can be gauged if μ_a is equivariant (31), in which case the gauged action is the (1,0) supersymmetrisation of (29), given by

$$S = \frac{1}{2} \int d^2x d\theta \psi_A \hat{D}\psi_A \quad (43)$$

where

$$\hat{D}\psi_A = \tilde{D}\psi_A + A^a \mu_a^{AB} \psi_B \quad (44)$$

REFERENCES

1. A. Sen, Phys. Rev. D3 (1985) 2162; Phys Rev Lett 55 (1985) 1846; C. Callan, D. Friedan, E. Martinec and M.J. Perry, Nucl. Phys.262 (1985) 593.
2. K. Bardacki, E. Rabinovici and B. S aring, Nucl. Phys.299 (1988) 157; H.J. Schnitzer, Nucl. Phys.324 (1989) 412; D. Karabali and H.J. Schnitzer, Nucl. Phys.329 (1990) 649; K. Gaw dzki and A. Kupiainen, Phys. Lett. 215B (1988) 119, Nucl. Phys. B320(FS) (1989) 625.
3. M. Ro ek and E. Verlinde, Nucl. Phys. **B373** (1992), 630.
4. X.C. de la Ossa and F. Quevedo, Nucl. Phys. **B403** (1993), 377.
5. C.M. Hull, G. Papadopoulos and P.K. Townsend, *Potentials for (p,0) and (1,1) supersymmetric sigma-models with torsion*, preprint DAMTP/R-93/8.
6. C.M. Hull and B. Spence, Phys. Lett. **B232** (1989), 204.
7. I. Jack, D.R.T. Jones, N. Mohammedi and H. Osborn, Nucl. Phys. **B332** (1990), 359.

8. C.M. Hull and B. Spence, Nucl. Phys. **B353** (1991), 379.
9. C.M. Hull, G. Papadopoulos and B. Spence, Nucl. Phys. **B363** (1991) 593.
10. C. M. Hull and E. Witten, Phys. Lett. 160B (1985) 398.
11. G. Papadopoulos, Phys. Lett. **B238** (1990), 75.
12. C.M. Hull, in preparation.
13. R. Brooks, F. Muhammad and S.J. Gates, Jr., Nucl. Phys. **268** (1986) 599.